




Distance between two points

Created: Thursday, 14 July 2011 07:49 | Published: Thursday, 14 July 2011 07:49 | Written by [Super User](#) | [Print](#)



The distance between any two points in the plane is the length of the line


segment joining them.

The distance between two points $P(x_1, y_1)$ and $Q(x_2, y_2)$ is given by

$$PQ = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

$$PQ = \sqrt{(\text{Difference of abscissae})^2 + (\text{Difference of ordinates})^2}$$

Let $P(x_1, y_1)$ and $Q(x_2, y_2)$ are two given points in the plane.



Draw PM and QL perpendicular from P and Q on x -axis. From P draw PN perpendicular to QL on x -axis. Then,

$$OM = x_1$$

$$OL = x_2$$

$$PM = y_1$$

$$QL = y_2$$

$$PN = ML = OL - OM = x_2 - x_1$$

$$QN = QL - NL = QL - PM = y_2 - y_1$$

So, $\triangle PNQ$ is a right triangle right angled at N . Therefore, by Pythagoras theorem, we have

$$PQ^2 = PN^2 + QN^2$$

$$PQ^2 = (x_2 - x_1)^2 + (y_2 - y_1)^2$$

$$PQ = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

Hence, distance between any two points is given by

$$\sqrt{(\text{Difference of abscissae})^2 + (\text{Difference of ordinates})^2}$$

Important Note:

- If O is the origin and $P(x, y)$ is any point, then from the above formula, we have

$$OP = \sqrt{(x - 0)^2 + (y - 0)^2}$$

$$OP = \sqrt{x^2 + y^2}$$

- In order to prove that a given figure is a:

i) Square, prove that four sides are equal and the diagonals are also equal.

- ii) Rhombus, prove that four sides are equal.
- iii) Rectangle, prove that opposite sides are equal and the diagonals are also equal.
- iv) Parallelogram, prove that opposite sides are equal.
- v) Parallelogram but not a rectangle, prove that its opposite sides are equal but the diagonals are not equal.
- vi) Rhombus but not a square, prove that its all sides are equal but the diagonals are not equal.
- For three points to be [collinear](#), prove that the sum of the distances between two pairs of points is equal to the third pair of points.

Example: Find the distance between the points: P (-6, 7) and Q (-1, -5)

Here $x_1 = -6$, $y_1 = 7$ and $x_2 = -1$, $y_2 = -5$

$$PQ = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

$$PQ = \sqrt{(-1 + 6)^2 + (-5 - 7)^2}$$

$$PQ = \sqrt{25 + 144}$$

$$PQ = \sqrt{169} = 13 \text{ units}$$

Example: Find a point on x – axis which is equidistant from A (2, -5) and B (-2, 9).

We know that a point on x – axis is of the form (x, 0). SO, let P(x, 0) be the point equidistant from A (2, -5) and B (-2, 9). Then,
 $PA = PB$

$$\sqrt{(x - 2)^2 + (0 + 5)^2} = \sqrt{(x + 2)^2 + (0 - 9)^2}$$

$$(x - 2)^2 + 25 = (x + 2)^2 + 81$$

$$x^2 - 4x + 4 + 25 = x^2 + 4x + 4 + 81$$

$$-8x = 56$$

$$x = -7$$

Hence, the required point is (-7, 0)

Now try it yourself! Should you still need any help, [click here](#) to schedule live online session with e Tutor!

About eAge Tutoring :

[eAgeTutor.com](#) is the premium online tutoring provider. Using materials developed by highly qualified educators and leading content developers, a team of top-notch software experts, and a group of passionate educators, eAgeTutor works to ensure the success and satisfaction of all of its students.

[Contact us](#) today to learn more about our guaranteed results and discuss how we can help make the dreams of the student in your life come true!

Reference Links :

- <http://en.wikipedia.org/wiki/Distance>
- <http://en.wikipedia.org/wiki/Abcissa>
- <http://www.thefreedictionary.com/ordinates>
- <http://en.wikipedia.org/wiki/Perpendicular>
- http://en.wikipedia.org/wiki/Pythagorean_theorem
- <http://www.thefreedictionary.com/collinear>

Category:ROOT

[Joomla SEF URLs by Artio](#)