#### **INDEFINITE INTEGRALS – II**

Created: Monday, 14 November 2011 10:40 | Published: Monday, 14 November 2011 10:40 | Written by Super User | Print

# **Integration by Substitution**

f(x) dx.

Here the given integral f(x) dx can be transformed into another form by changing the <u>independent variable</u> 'x' to 't' by substituting x=g(t).

Consider I = ?f(x) dxPut x=g(t) so that dx/dt = g'(t)We can write dx = g'(t)dtHence I = ?f(x) dx = ?f[g(t)] g'(t)dtThis change of variable formula is one of the important tools available to us in the name of integration by substitution. Usually, we make a substitution for a function whose derivative also occurs in the integrand. We can have a glance at few examples.  $\int \tan x \, dx = \int \sin x \, dx$ Cosx Put cosx = t, so that -sinxdx=dt ∫tanx dx = ∫-dt i) t  $= - \log|t| + C$  $= -\log|\cos x| + C$  $= \log |\cos x|^{-1} + C$  $= \log |1/\cos x| + C$  $= \log|secx| + C$ Hence  $2 \tan x \, dx = \log |\sec x| + C$ Similarly we can find the integral of cotx also  $2 \cot x \, dx = \log |\sin x| + C$ ∫(logx)<sup>2</sup> dx ii) X Here we know the derivative of logx is 1/x, so put logx = t  $\log x = t$ 1/x dx = dt2 ∫(logx)<sup>2</sup> dx = ?t dt X  $=t^{3}/3 + C$ (logx)<sup>3</sup> 

Using substitution technique, we can find the following standard integrals.

```
i) 2 \tan x \, dx = \log |\sec x| + C
```

```
ii) 2\cot x \, dx = \log |\sin x| + C
```

```
iii) \operatorname{?secx} dx = \log |\operatorname{secx} + \tan x| + C
```

```
iv) 2 \operatorname{cosecx} dx = \log |\operatorname{cosecx} - \operatorname{cotx}| + C
```

### Integration using trigonometric identities

When the integrand involves some trigonometric functions, we use some well known identities to find the integrals. Most commonly used <u>trigonometric identities</u> are,

- $\cos^2 x = [1 + \cos 2x]/2$
- $\sin^2 x = [1 \cos 2x]/2$
- $2\cos(x+y) + \cos(x-y)$
- $2\sin x \sin y = \cos(x-y) \cos(x+y)$
- $2\sin x \cos y = \sin(x+y) + \sin(x-y)$
- $2\cos x \sin y = \sin(x+y) \sin(x-y)$
- $\sin^3 x = [3\sin x \sin 3x]/4$
- $\cos^3 x = [3\cos x + \cos 3x]/4$

## Integrals of special form

While integrating functions of the form  $\int px+q \, dx$  and  $\int px+q \, dx$  $ax^2+bx+c$   $\sqrt{ax^2+bx+c}$ 

where p, q, a, b and c are constants we have to use another method.

Assume that px+q = A d  $(ax^2+bx+c) + B = A(2ax+b) + B$ 

dx

Which means that Numerator = A derivative  $(ax^2+bx+c) + B$ 

To determine A and B, we equate from both sides the coefficients of 'x' and constant terms. After getting the values of A and B we can reduce the integral to one of the known forms.

### **Integration by Parts**

Integration by Parts is applicable only for the functions which are expressed as product of functions.

If 'u' and 'v' are any two differentiable functions of a single variable 'x', then

uv dx = uvdx - v(uvdx) dx

OR

f(x) g(x)dx = f(x) g(x)dx - f(x)g(x)dx]dx.

This formula can stated as follows: "The integral of the product of two functions = (first function) x (integral of the second function) – Integral of [(derivative of first function) x (integral of the second function)]".

Example: Find ?x sin3x dx

Solution:  $x \sin 3x \, dx = x \sin 3x \, dx - 2[d(x) \sin 3x \, dx] \, dx$ 

 $dx = x (-\cos 3x)/3 - ?1 (-\cos 3x)/3 dx$ 

 $-x\cos 3x + \sin 3x + C$ 

**Note:** While performing Integration by Parts we have memorize "ILATE" which gives the order in which the functions are to be taken. Inverse and <u>logarithmic functions</u> must be taken as first function since they doesn't have integrals.

I = Inverse function

L = Logarithmic function

A = Algebraic Function

T = Trigonometric function

E = Exponential function

# Integrals of the type $e^{x} [f(x) + f'(x)] dx$

We have I=?e<sup>x</sup> [f(x) + f'(x)]dx = ?e<sup>x</sup> f(x)dx + ?e<sup>x</sup> f'(x)dx = I1 + ?e<sup>x</sup> f'(x) dx where I1=?e<sup>x</sup> f(x) dx .....(1) I1 = ?e<sup>x</sup> f(x) dx = ?f(x) e<sup>x</sup> dx [Take f(x) as 1st function] = f(x) e<sup>x</sup> - ?f'(x) e<sup>x</sup> dx = f(x) e<sup>x</sup> - ?f'(x) e<sup>x</sup> dx + C (1) becomes, I = e<sup>x</sup> f(x) - ?f'(x) dx + ?f'(x) dx + C = e<sup>x</sup> f(x) + C Hence ?e<sup>x</sup> [f(x) + f'(x)] dx = e<sup>x</sup> f(x) + C Example: Find ?e<sup>x</sup> [sinx + cosx] dx Solution: ?e<sup>x</sup> [sinx + cosx] dx = e<sup>x</sup> sinx + C [If f(x)=sinx then f'(x)=cosx]

### Integrals of some more types

Here we discuss some special types of standard integrals based on the technique of integration by parts:

i) 
$$\int \sqrt{x^2 - a^2} \, dx = x\sqrt{x^2 - a^2} - \frac{a^2 \log |x + \sqrt{x^2 - a^2}| + C}{2}$$
  
ii)  $\int \sqrt{x^2 + a^2} \, dx = x\sqrt{x^2 + a^2} + \frac{a^2 \log |x + \sqrt{x^2 + a^2}| + C}{2}$   
iii)  $\int \sqrt{a^2 - x^2} \, dx = x\sqrt{a^2 - x^2} + \frac{a^2 \sin^{-1}}{2} \left(\frac{x}{a}\right) + C$ 

Now try it yourself! Should you still need any help, click here to schedule live online session with e Tutor!

#### **About eAge Tutoring:**

<u>eAgeTutor.com</u> is the premium online tutoring provider. Using materials developed by highly qualified educators and leading content developers, a team of top-notch software experts, and a group of passionate educators, eAgeTutor works to ensure the success and satisfaction of all of its students.

<u>Contact us</u> today to learn more about our tutoring programs and discuss how we can help make the dreams of the student in your life come true!

#### **Reference Links:**

- <u>http://en.wikipedia.org/wiki/Integration\_by\_parts</u> http://www.analyzemath.com/logfunction/logfunction.html
- http://en.wikipedia.org/wiki/Integration\_by\_substitution
- http://en.wikipedia.org/wiki/List\_of\_trigonometric\_identities
  http://en.wikipedia.org/wiki/Dependent\_and\_independent\_variables

Category:ROOT