


# Refraction

Created: Thursday, 25 August 2011 04:53 | Published: Thursday, 25 August 2011 04:53 | Written by [Super User](#) | [Print](#)

## Introduction to Refraction


The light rays from the straw are refracted (or bent) when they cross from water back into the air before reaching your eyes



### What is refraction?

When a ray of light passes from one medium to another, the direction of the ray changes. This is known as refraction. The refracted ray will bend toward the normal line when the second medium is denser than the first medium and vice versa. To learn about the normal line, read the eAgeTutor article on “reflection.”

How much a ray bends depends on the [indices of refraction](#) of the two media, and can be described quantitatively by [Snell's Law](#). The speed of light is reduced in the slower medium, and the wavelength is shortened proportionately. The frequency is unchanged, however, since frequency is a characteristic of the source of the light and is unaffected by medium changes. Image formation by lenses and the eye requires refraction.



### The Cause of Refraction

When a light wave encounters a new medium or an obstacle in its path, it can respond in various ways, and refraction is merely one of several of these possible boundary behaviors. A change in both the speed and [wavelength](#) of a light wave accompanies the transmission of light across a boundary between two media. The light wave will not only change directions at the boundary, but will also speed up or slow down and transform itself into a wave with a larger or a shorter wavelength. The only time that a wave can be transmitted across a boundary, change its speed, but not refract is when the light wave approaches a boundary from a direction that is perpendicular to the boundary. Whenever a light wave changes speed and approaches a boundary at an angle,

refraction will be observed.

## Laws of Refraction

### There are two laws of refraction

Law 1. The [incident ray](#), the normal line, and the refracted ray all lie in the same plane.

Law 2. The ratio:  $\frac{\text{sine of the angle of incidence (i)}}{\text{sine of the angle of refraction (r)}}$

is constant for light of a given color (i.e., [frequency](#)) crossing the boundary of two given media. This ratio is called the refractive index (n).

$$n = \frac{\sin i}{\sin r}$$

The refractive index of a transparent substance is measured when light enters the substance from a vacuum, but for all practical purposes we can take it as light entering the substance from air. The greater the value of the [refractive index](#), the more the light will bend. [Refraction](#) is caused by the fact that light travels at different speeds in different media; it can also be shown that the refractive index is equal to the ratio of the speeds in the two media.

Want to know more about refraction? [Click here](#) to schedule a live session with an eAge eTutor!

## About eAge Tutoring:

[eAgeTutor.com](#) is the premium online tutoring provider. Using materials developed by highly qualified educators and leading content developers, a team of top-notch software experts, and a group of passionate educators, eAgeTutor works to ensure the success and satisfaction of all of its students.

[Contact us](#) today to learn more about our guaranteed results and discuss how we can help make the dreams of the student in your life come true!

## Reference links:

- <http://www.ncsu.edu/scivis/lessons/reflection/reflection/tsld>
- [http://www.ence.kosmix.com/topic/Angle\\_of\\_incidence](http://www.ence.kosmix.com/topic/Angle_of_incidence)
- [http://www.absoluteastronomy.com/topics/Refractive\\_index](http://www.absoluteastronomy.com/topics/Refractive_index)
- <http://www.answers.com/topic/frequency>

Category:ROOT

[Joomla SEF URLs by Artio](#)