


Monatomic gases



Figuring out the degree of freedom for [monatomic gases](#) is simple because



[internal energy](#) is stored only in the translational motion of the atoms. That is, the atoms move as a point-like body and can move in three independent directions.

For a monatomic gas, there are three translational degrees of freedom corresponding to the three axes of movement.

As its rotational moment of [inertia](#) is very low (due to small atomic radius), the degree of freedom associated with rotation does not make an appreciable contribution to the kinetic energy. Therefore, a monatomic gas has 3 degrees of freedom.

Degree of freedom $f = 3$ (all translational)

Diatomic gases



For diatomic and polyatomic molecules, rotational


motion is significant.

In these cases both translational and rotational motion can contribute to [the heat capacity](#):

For a diatomic gas, in addition to the three degrees for [translatory motion](#), there are two degrees of rotatory motion around axes perpendicular to the bond between the atoms.

Degree of freedom, $f = 5$ (3 translational + 2 rotational) at room temperature $f = 7$ (3 translational + 2 rotational + 2 vibrational) at high temperature

Polyatomic gas



A polyatomic (non-linear) molecule (such as NH_3) can rotate about any of three

co-ordinate axes. Hence it has 6 degrees of freedom, 3 translational and 3 rotational. At room temperature, [apolyatomic gas](#) molecule has vibrational energy greater than that of a diatomic gas. The atoms within the molecule may also vibrate with respect to each other at high temperature. In such cases, the molecule will have an additional degree of freedom, due to vibrational motion. The vibrational motion of a molecule can contribute as much as to two further degrees of freedom. Vibrational motion is not allowed with diatomic molecules but can occur with polyatomic molecules

Degree of freedom, $f = 6$ (3 translational + 3 rotational) at room temperature

$f = 8$ (3 translational + 3 rotational + 2 vibrational) at high temperature

Want to know more about degrees of freedom? [Click here](#) to schedule a live session with an eAge eTutor!

About eAge online Tutoring

[eAgeTutor.com](#) is the premium online tutoring provider. Using materials developed by highly qualified educators and leading content developers, a team of top-notch software experts, and a group of passionate educators, eAgeTutor works to ensure the success and satisfaction of all of its students.

[Contact us](#) today to learn more about our guaranteed results and discuss how we can help make the dreams of the student in your life come true!

Reference links

- http://www.en.wikipedia.org/wiki/Kinetic_energy
- <http://www.en.wikipedia.org/wiki/temperature>
- <http://www.hyperphysics.phy-astr.gsu.edu/hbase/thermo/inteng.html>
- <http://www.chm.davidson.edu/vce/calorimetry/heatcapacity.html>

Category:ROOT

[Joomla SEF URLs by Artio](#)