

INDEFINITE INTEGRALS – I

Created: Monday, 14 November 2011 09:15 | Published: Monday, 14 November 2011 09:15 | Written by [Super User](#) | [Print](#)

Methods of Integration

$$\int f(x) dx.$$

We have already discussed that [integrals](#) of some functions are obtained from their derivatives. It was based on inspection which means that on search of a function F whose derivative is 'f'. Method of inspection is not suitable for many of the functions. So we were forced to find additional methods to reduce a function to its standard form. The methods which we follow are

- 1) Integration using [Partial Fractions](#)
- 2) Integration by substitution
- 3) Integration by Parts.

Some Standard Integrals

- 1) $\int x^n dx = \frac{x^{n+1}}{n+1} + C, n \neq -1$
- 2) $\int dx = x + C$
- 3) $\int \cos x dx = \sin x + C$
- 4) $\int \sin x dx = -\cos x + C$
- 5) $\int \sec^2 x dx = \tan x + C$
- 6) $\int \operatorname{cosec}^2 x dx = -\cot x + C$
- 7) $\int \sec x \tan x dx = \sec x + C$
- 8) $\int \operatorname{cosec} x \cot x dx = -\operatorname{cosec} x + C$
- 9) $\int e^x dx = e^x + C$
- 10) $\int 1/x dx = \log|x| + C$
- 11) $\int dx / \sqrt{1-x^2} = \sin^{-1} x + C$
- 12) $\int dx / \sqrt{1-x^2} = -\cos^{-1} x + C$
- 13) $\int dx / (1+x^2) = \tan^{-1} x + C$
- 14) $\int dx / (1+x^2) = -\cot^{-1} x + C$
- 15) $\int dx / (x^2-1) = \sec^{-1} x + C$

Integration using Partial Fractions

We have already learned that a [rational function](#) is defined as the ratio of two polynomials in the form P(x)/Q(x), where P(x) and Q(x) are polynomials in 'x' and Q(x) ≠ 0. If the degree of P(x) is less than the degree of Q(x), then it is called proper otherwise improper.

The improper rational functions can be reduced to the proper rational functions by long division process. It can be expressed in the form

$$\frac{P(x)}{Q(x)} = T(x) + \frac{P_1(x)}{Q(x)}$$

where $T(x)$ is a polynomial in 'x' and $P_1(x)/Q(x)$ is a proper rational function. Here the rational functions which we consider for integration purpose will be those whose denominators can be factorized into linear and quadratic factors.

To evaluate $\int P(x) dx$, where $P(x)$ is a proper rational function

$$\frac{P(x)}{Q(x)}$$

we can express the integrand as a sum of simpler rational functions which is called the method of [partial fractions](#). After decomposing into simpler rational functions we can integrate it easily.

Different Forms of Partial Fractions

The following table indicates the types of simpler partial fractions that are to be associated with various kinds of rational functions. Before decomposing the rational functions we have to determine which form of partial fraction is to be applied. Here A, B and C are constants (real numbers) to be determined by suitable method.

Sl. No:	Form of rational function	Form of the partial fraction
1	$\frac{px+q}{(x-a)(x-b)}, a \neq b$	$\frac{A}{x-a} + \frac{B}{x-b}$
2	$\frac{px+q}{(x-a)^2}$	$\frac{A}{x-a} + \frac{B}{(x-a)^2}$
3	$\frac{px^2+qx+r}{(x-a)(x-b)(x-c)}$	$\frac{A}{x-a} + \frac{B}{x-b} + \frac{C}{x-c}$
4	$\frac{px^2+qx+r}{(x-a)^2(x-b)}$	$\frac{A}{x-a} + \frac{B}{(x-a)^2} + \frac{C}{x-b}$
5	$\frac{px^2+qx+r}{(x-a)(x^2+bx+c)}$	$\frac{A}{x-a} + \frac{Bx+C}{x^2+bx+c}$

where x^2+bx+c cannot be factorized.

Find $\int \frac{2x}{x^2+3x+2} dx$

$$\int \frac{2x}{x^2+3x+2} dx = \int \frac{2x}{(x+2)(x+1)} dx$$

Example: Take $\frac{2x}{(x+2)(x+1)} = \frac{A}{x+2} + \frac{B}{x+1}$ (1)

$$2x = A(x+1) + B(x+2) \quad [\text{Taking L.C.M}]$$

For finding A and B we have to assign values to 'x'.

When $x=-1$, we get $B=-2$

When $x=-2$, we get $A=4$

(1) becomes

$$\frac{2x}{(x+2)(x+1)} = \frac{4}{(x+2)} + \frac{-2}{(x+1)}$$

$$\int \frac{2x}{(x+2)(x+1)} dx = \int \frac{4}{(x+2)} dx - \int \frac{2}{(x+1)} dx$$

$$= 4 \log(x+2) - 2 \log(x+1) + C$$

Integrals of some special function

When we come across the integrals like $\int \frac{dx}{ax^2+bx+c}$ and $\int \frac{dx}{\sqrt{ax^2+bx+c}}$ where the

denominator cannot be factorized as linear factors; we have to go for some other method. For solving such integrals we have to learn six formulas from which we get the answer easily.

The formulas are given below.

$$1) \int \frac{dx}{x^2-a^2} = \frac{1}{2a} \log \left| \frac{x-a}{x+a} \right| + C$$

$$2) \int \frac{dx}{a^2-x^2} = \frac{1}{2a} \log \left| \frac{a+x}{a-x} \right| + C$$

$$3) \int \frac{dx}{x^2+a^2} = \frac{1}{a} \tan^{-1} \left(\frac{x}{a} \right) + C$$

$$4) \int \frac{dx}{\sqrt{x^2-a^2}} = \log | x + \sqrt{x^2-a^2} | + C$$

$$5) \int \frac{dx}{\sqrt{a^2-x^2}} = \sin^{-1} (x/a) + C$$

$$6) \int \frac{dx}{\sqrt{x^2+a^2}} = \log | x + \sqrt{x^2+a^2} | + C$$

The remaining two methods of integration will be discussed in the article named "Indefinite Integrals – II".

Now try it yourself! Should you still need any help, [click here](#) to schedule live online session with e Tutor!

About eAge Tutoring:

eAgeTutor.com is the premium online tutoring provider. Using materials developed by highly qualified educators and leading content developers, a team of top-notch software experts, and a group of passionate educators, eAgeTutor works to ensure the success and satisfaction of all of its students.

[Contact us](#) today to learn more about our tutoring programs and discuss how we can help make the dreams of the student in your life come true!

Reference Links:

- http://en.wikipedia.org/wiki/Partial_fractions_in_integration

- http://en.wikipedia.org/wiki/Rational_function
- http://en.wikipedia.org/wiki/Lists_of_integrals
- <http://www.sosmath.com/algebra/pfrac/pfrac07.html>

Category:ROOT

[Joomla SEF URLs by Artio](#)