


RIEMANN SUM AND TRAPEZOIDAL RULE

Created: Friday, 18 November 2011 09:42 | Published: Friday, 18 November 2011 09:42 | Written by [Super User](#) | [Print](#)


Riemann Sum Approximation



The definite integral of a continuous function 'f' over an interval [a, b] is computed as $\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{k=1}^n f(x_k^*) \Delta x$, where the sum that appears on the right side is called [Riemann sum](#). In this formula, the interval [a, b] is divided into n subintervals of width $\Delta x = (b-a)/n$, and x_k^* denotes an arbitrary point in the k^{th} sub-interval. It follows that as n increases the Riemann sum will eventually be a good approximation to the integral, which we denote by writing

$$\int_a^b f(x) dx \approx \sum_{k=1}^n f(x_k^*) \Delta x$$

$$\int_a^b f(x) dx \approx \Delta x [f(x_1^*) + f(x_2^*) + \dots + f(x_n^*)]$$




Here we denote the values of 'f' at the endpoints of the subintervals by

$y_0=f(a)$, $y_1=f(x_1)$, $y_2=f(x_2)$, ..., $y_{n-1}=f(x_{n-1})$, $y_n=f(b)$ and we will denote the values of f at the midpoints of the subintervals by y_{m1} , y_{m2} , ..., y_{mn}

Trapezoidal Approximation

The left-hand and right hand endpoint approximations are rarely used in applications; however, if we take the average of the left-hand and right hand endpoint approximations, we obtain a result, called the trapezoidal approximation, which is commonly used as,

$$\int_a^b f(x) dx \approx (b-a)/2n [y_0 + 2y_1 + \dots + 2y_{n-1} + y_n]$$



represents the area under $f(x)$ over $[a, b]$. Geometrically, the trapezoidal approximation formula results if we approximate this area by the sum of the trapezoidal areas as shown in the figure

Left end point Approximation: The formula for evaluating left end point approximation is given by

$$\int_a^b f(x)dx = (b-a)/n [y_0 + y_1 + \dots + y_{n-1}]$$

Right Endpoint Approximation: The formula for evaluating right end point approximation is given by

$$\int_a^b f(x)dx = (b-a)/n [y_1 + y_2 + \dots + y_n]$$

Mid-Point Approximation: The formula for evaluating midpoint approximation is given by

$$\int_a^b f(x)dx = (b-a)/n [y_{m1} + y_{m2} + \dots + y_{mn}] \text{ where } m_1, m_2, \dots, m_n \text{ represents the mid values.}$$

Example: Use Trapezoidal rule to approximate $\int_0^{\pi/2} \sin x \, dx$ using $n=10$ sub intervals

Solution: $a=0$, $b=\pi/2$, $n=10$ and $f(x)=\sin x$, $(b-a)/n = \pi/20$

i	0	1	2	3	10
x_i	0	$\pi/10$	$2\pi/10$	$3\pi/10$	$10\pi/10$
y_i	$\sin(0)$	$\sin(\pi/10)$	$\sin(2\pi/10)$	$\sin(3\pi/10)$	$\sin(10\pi/10)$

$$\begin{aligned} \int_0^{\pi/2} \sin x \, dx &= (\pi/20)[y_0 + 2y_1 + 2y_2 + \dots + 2y_{n-1} + y_n] \\ &= (\pi/20) [\sin(0) + 2\sin(\pi/10) + 2\sin(2\pi/10) + \dots + \sin(10\pi/10)] \\ &= 1.983523538 \end{aligned}$$

Comparison of the Midpoint and Trapezoidal Approximations

The table below shows the comparison between midpoint and trapezoidal approximations for the function $\ln 2 = \int_1^2 (1/x)dx$ with $n=10$ subdivisions

Midpoint Approximation

i	Midpoint (m_i)	$y_{mi} = f(m_i) = 1/m_i$
1	1.05	0.952380952
2	1.15	0.869565217
3	1.25	0.800000000
4	1.35	0.740740741
5	1.45	0.689655172
6	1.55	0.645161290
7	1.65	0.606060606
8	1.75	0.571428571
9	1.85	0.540540541
10	1.95	0.512820513
		<u>6.928353603</u>

$$\int_1^2 (1/x)dx = (0.1)(6.928353603) = 0.692835360$$

Trapezoidal Approximation

i	Endpoint(x_i)	$y_i=f(x_i)=1/x_i$	Multiplier (w_i)	$w_i y_i$
0	1.0	1.000000000	1	1.000000000
1	1.1	0.909090909	2	1.818181818
2	1.2	0.833333333	2	1.666666667
3	1.3	0.769230769	2	1.538461538
4	1.4	0.714285714	2	1.428571429
5	1.5	0.666666667	2	1.333333333
6	1.6	0.625000000	2	1.250000000
7	1.7	0.588235294	2	1.176470588
8	1.8	0.555555556	2	1.111111111
9	1.9	0.526315789	2	1.052631579
10	2.0	0.500000000	1	0.500000000
				13.875428063

$$\int_1^2 (1/x)dx = (0.05)(13.875428063) = 0.693771403$$

The value of $\ln 2$ is rounded to nine decimal places and we have seen that [midpoint approximation](#) produces a more accurate result than the trapezoidal approximation. Hence we can conclude that,

If f be a continuous on $[a, b]$ and let $|E_M|$ and $|E_T|$ be the absolute errors that result from the midpoint and trapezoidal approximations of $\int_a^b f(x)dx$ using n subintervals.

a) If the graph of f is either concave up or concave down on (a, b) , then $|E_M| < |E_T|$, which means that the error from the midpoint approximation is less than from the trapezoidal approximation.

b) If the graph of ' f ' is concave down on (a, b) then $T_n < \int_a^b f(x)dx < M_n$

c) If the graph of ' f ' is concave up on (a,b) , then $M_n < \int_a^b f(x)dx < T_n$

Simpson's Rule

[Simpson's Rule](#) is given by

$$S_{2n} = \frac{1}{3} (2M_n + T_n)$$

$$= \frac{1}{3} \left(\frac{b-a}{2n} \right) [y_0 + 4y_1 + 2y_2 + 4y_3 + 2y_4 + \dots + 2y_{n-2} + 4y_{n-1} + y_{2n}]$$

$$\text{Where } M_n = \left(\frac{b-a}{2n} \right) [2y_1 + 2y_3 + \dots + 2y_{n-1}]$$

$$T_n = \left(\frac{b-a}{2n} \right) [y_0 + 2y_2 + 2y_4 + \dots + 2y_{n-2} + y_{2n}]$$

$$\text{Also } \int_a^b f(x) dx \approx S_{2n}$$

We denote the absolute error in this approximation by

$$|E_S| = \left| \int_a^b f(x) dx - S_{2n} \right|$$

Now try it yourself! Should you still need any help, [click here](#) to schedule live online session with e Tutor!

About eAge Tutoring:

eAgeTutor.com is the premium online tutoring provider. Using materials developed by highly qualified educators and leading content developers, a team of top-notch software experts, and a group of passionate educators, eAgeTutor works to ensure the success and satisfaction of all of its students.

[Contact us](#) today to learn more about our tutoring programs and discuss how we can help make the dreams of the student in your life come true!

Reference links:

- http://en.wikipedia.org/wiki/Riemann_sum
- http://en.wikipedia.org/wiki/Trapezoidal_rule
- http://en.wikipedia.org/wiki/Simpson's_rule
- <http://www.purplemath.com/modules/numeric.htm>

Category:ROOT

[Joomla SEF URLs by Artio](#)