Buy from flipkart.com

INVERTIBLE MATRICES

Created: Friday, 25 November 2011 06:11 | Published: Friday, 25 November 2011 06:11 | Written by Super User | Print

What are Invertible matrices?

$$A^{-1} = \frac{1}{|A|}\begin{bmatrix} A_{11} & A_{21} & A_{31} \\ A_{12} & A_{22} & A_{32} \\ A_{13} & A_{32} & A_{33} \end{bmatrix}$$

If A is a square matrix of order m, and if there exists another square matrix B of the same order m such that AB = BA = I, then B is called the inverse matrix of A and is denoted by A^{-1}. In this case we say A is invertible.

Important Remarks:

- Inverse of a square matrix, if it exists, is unique.
- If A and B are invertible matrices of the same order, then $(AB)^{-1} = B^{-1}A^{-1}$

Inverse of a matrix by elementary operations

There are six operations (transformations) on a matrix, three of which are due to rows and three due to columns, which are known as elementary operations or transformations.

i) The interchange of any two rows or two columns. Symbolically the interchange of i^{th} and j^{th} rows is denoted by $R_i \leftrightarrow R_j$ and the interchange of i^{th} and j^{th} column is denoted by $C_i \leftrightarrow C_j$

ii) The multiplication of the elements of any row or column by a non zero number. Symbolically the multiplication of each element of the i^{th} row by k, where k?0 is denoted by $R_i \leftrightarrow kR_i$. The corresponding column operation is denoted by $C_i \leftrightarrow kC_i$

iii) The addition to the elements of any row or column, the corresponding elements of any other row or column multiplied by any non zero number. Symbolically, the addition to the elements of ith row, the corresponding elements of jth row multiplied by k is denoted by $R_i \leftrightarrow R_i + kR_j$ The corresponding column operation is denoted by $C_i \leftrightarrow C_i + kC_j$

Example: Obtain the inverse of the following matrix using elementary operation

$$A = \begin{pmatrix} 0 & 1 & 2 \\ 1 & 2 & 3 \\ 3 & 1 & 1 \end{pmatrix}$$

Solution: We know A=IA

$$\begin{pmatrix}0&1&2\\1&2&3\\3&1&1\end{pmatrix} = \begin{pmatrix}1&0&0\\0&1&0\\0&0&1\end{pmatrix} A$$

$$\begin{pmatrix}1&2&3\\0&1&2\\3&1&1\end{pmatrix} = \begin{pmatrix}0&1&0\\1&0&0\\0&0&1\end{pmatrix} A \qquad R_1 \longleftrightarrow R_2$$

$$\begin{pmatrix}1&2&3\\0&1&2\\0&-5&-8\end{pmatrix} = \begin{pmatrix}0&1&0\\1&0&0\\0&-3&1\end{pmatrix} A \qquad R_3 \rightarrow R_1-3R_1$$

$$\begin{pmatrix}1&0&-1\\0&1&2\\0&-5&-8\end{pmatrix} = \begin{pmatrix}-2&1&0\\1&0&0\\0&-3&1\end{pmatrix} A \qquad R_1 \rightarrow R_1-2R_2$$

$$\begin{pmatrix}1&0&-1\\0&1&2\\0&0&2\end{pmatrix} = \begin{pmatrix}-2&1&0\\1&0&0\\5&-3&1\end{pmatrix} A \qquad R_3 \rightarrow R_3+5R_2$$

$$\begin{pmatrix}1&0&-1\\0&1&2\\0&0&1\end{pmatrix} = \begin{pmatrix}-2&1&0\\1&0&0\\5/2&-3/2&½\end{pmatrix} A \qquad R_3 \rightarrow ½\ R_3$$

$$\begin{pmatrix}1&0&0\\0&1&2\\0&0&1\end{pmatrix} = \begin{pmatrix}½&-1/2&½\\1&0&0\\5/2&-3/2&½\end{pmatrix} A \qquad R_1 \rightarrow R_1+R_2$$

$$\begin{pmatrix}1&0&0\\0&1&0\\0&0&1\end{pmatrix} = \begin{pmatrix}½&-1/2&½\\-4&3&-1\\5/2&-3/2&½\end{pmatrix} A \qquad R_2 \rightarrow R_2-2R_3$$

$$\text{Hence } A^{-1} = \begin{pmatrix}½&-1/2&½\\-4&3&-1\\5/2&-3/2&½\end{pmatrix}$$

Now try it yourself! Should you still need any help,click here to schedule live online session with e Tutor!

About eAge Tutoring:

eAgeTutor.com is the premium online tutoring provider. Using materials developed by highly qualified educators and leading content developers, a team of top-notch software experts, and a group of passionate educators, eAgeTutor works to ensure the

success and satisfaction of all of its students.

Contact us today to learn more about our tutoring programs and discuss how we can help make the dreams of the student in your life come true!

Reference Links:

http://www.britannica.com/EBchecked/topic/561660/square-matrix

http://en.wikipedia.org/wiki/Invertible_matrix

http://www.purplemath.com/modules/mtrxrows.htm

Category:ROOT

Joomla SEF URLs by Artio